

## Acute pneumonia associated with Chlamydia pneumoniae infection : a case report

Sasithorn Likitnukul\*

Pongpun Nunthapisud\*\* Somying Tumwasorn\*\*

Likitnukul S, Nunthapisud P, Tumwasorn S. Acute pneumonia associated with Chlamydia pneumoniae infection : a case report. Chula Med J 1999 Dec; 43(12): 893-9

*Chlamydia pneumoniae is the recently recognized third species of Chlamydia associated with human disease. It is an important cause of respiratory tract infection. The disease occurs worldwide, more commonly in adults than in children. We report the case of a 12-year-old boy presented with acute respiratory symptoms. Roentgenogram of the chest showed minimal reticular infiltration in the right lower lung with minimal right pleural effusion confirming the diagnosis of pneumonia. The evidence of current C. pneumoniae infection was confirmed by the microimmunofluorescence test for C. pneumoniae IgM 1:64 , IgG 1:256 in acute serum., and IgM 1:128, IgG 1:256 in the second serum sample at a four-week interval. He received roxithromycin 150 mg twice a day for 14 days showing a good response. One should consider C. pneumoniae as a cause of acute community acquired pneumonia in children.*

**Key words:** *C. pneumoniae pneumonia, Pleural effusion.*

Reprint request: Likitnukul S, Department of Pediatrics, Faculty of Medicine, Chulalongkorn University, Bangkok 10330, Thailand.

Received for publication. September 1, 1999.

\*Department of Pediatrics, Faculty of Medicine, Chulalongkorn University

\*\*Department of Microbiology, Faculty of Medicine, Chulalongkorn University

ศศิธร ลิขิตบุญกุล, ผ่องพรรณ นันทากสิสุทธิ์, สมหญิง ธัมวาสร. ปอดบวมจากการติดเชื้อคลาไมเดีย นิวโมเนีย : รายงานผู้ป่วย 1 ราย. จุฬาลงกรณ์เวชสาร 2542 ธ.ค; 43(12): 893-9

คลาไมเดีย นิวโมเนีย จัดเป็นคลาไมเดียชนิดที่สามก่อโรคในคน โดยเป็นสาเหตุสำคัญของโรคติดเชื้อทางเดินหายใจ การติดเชื้อนี้พบได้ทั่วโลกพบในผู้ใหญ่มากกว่าเด็ก บทความนี้เป็นรายงานผู้ป่วยเด็กชาย 1 ราย อายุ 12 ปี มีอาการของโรคติดเชื้อทางเดินหายใจ การตรวจภาพรังสีปอดพบความผิดปกติบริเวณชายปอดด้านขวาเป็นลักษณะ : *reticular infiltration* และ *minimal pleural effusion*. ผลการตรวจโดยวิธีไมโครอิมมูโนฟลูออเรสเซน แสดงว่ากำลังมีการติดเชื้อคลาไมเดีย นิวโมเนีย ในการตรวจน้ำเหลืองครั้งแรกพบระดับ IgM 1 : 64, IgG : 256 และครั้งที่สองห่างกัน 4 สัปดาห์พบระดับ IgM 1 : 128 และ IgG 1 : 256 ผู้ป่วยตอบสนองดีต่อการรับประทานยา *roxithromycin* ขนาด 150 มก. วันละ 2 ครั้ง เป็นเวลา 14 วัน แพทย์ควรคำนึงถึงเชื้อคลาไมเดีย นิวโมเนีย ด้วยว่าอาจเป็นสาเหตุของโรคปอดบวมในเด็ก

*Chlamydia pneumoniae* is a recently recognized human pathogen that can cause pneumonia, bronchitis, pharyngitis, otitis media, sinusitis,<sup>(1)</sup> and may also trigger acute episodes of wheezing in patients with asthma.<sup>(2)</sup> In adults, there have been several reports of an association between coronary artery disease, atherosclerotic syndromes and *C. pneumoniae* infection.<sup>(3-5)</sup> Worldwide seroepidemiological studies revealed that approximately 50% of adults have antibodies to this pathogen. The prevalence undergoes the greatest change in children between 5 and 15 years of age and continues to rise throughout adult life reaching 70-80% in the elderly.<sup>(6)</sup> The population prevalence was higher in tropical, less developed countries than in more northern developed countries. Nunthapisud et al.<sup>(7)</sup> reported a seroprevalence of 37% in Thai school children on a worldwide scale. *C. pneumoniae* has been associated with pneumonia in 6-19%.<sup>(8-15)</sup> As yet, there has not been any study of this infection in connection with lower respiratory tract infection in Thai children. We herein report a child with pneumonia associated with *C. pneumoniae* infection.

#### Case Report

A 12-year-old boy presented with sore throat and a cough producing green mucoid sputum for seven days. He reported a high fever for the previous three days. Physical examination revealed a body temperature of 39°C, respiratory rate of 24 beats/min, pulse rate of 100 beats/min, and blood pressure of 90/50 mm Hg. The pharynx was erythematous, and medium crepitation was heard at the base of the right lung. Initial laboratory investigations included a white blood cell count of 15,300/mm<sup>3</sup> with 81% neutrophils,

9% lymphocytes, 2% atypical lymphocytes, 8% monocytes. The hematocrit was 34% and the platelet count 345,000.

A chest roentgenogram showed minimal reticular infiltration in the right lower lung with minimal right pleural effusion. (Fig. 1) A diagnosis of pneumonia, probably due to mycoplasma, was made, and the patient was treated with roxithromycin 150 mg twice a day. Throat swab was performed for polymerase chain reaction (PCR) detection of *Mycoplasma pneumoniae*, and blood was drawn for serology for *M. pneumoniae* and *Chlamydia pneumoniae*. When seen in the clinic seven days later, the patient reported




Figure 1. Chest radiograph demonstrates minimal reticular infiltration in the right lower lung with minimal right pleural effusion.

improvement beginning two days after having started the treatment with roxithromycin. The fever had disappeared on the third day of treatment but there was still some coughing. On examination, the temperature and breath sounds were normal. He received roxithromycin to complete 14 days of treatment, which resulted in complete recovery. A follow up chest radiograph four weeks later showed complete clearing of infiltration and pleural effusion in the right lower lung. The second blood specimens were sent for antibody testing for *M. pneumoniae* and *C. pneumoniae*.

Laboratory results included negative PCR for *M. pneumoniae*, negative antibody for *M. pneumoniae* by particle agglutination test with a titer below 1:40 with both the first and second specimen. Microimmunofluorescence for *C. pneumoniae* with the first serum showed IgM 1:64 and IgG 1:256, and the second serum sample four weeks later showed IgM 1:128, IgG 1:256. Subsequently he was lost to follow-up.

## Discussion

Acute *C. pneumoniae* infection can range from asymptomatic to life - threatening. Pneumonia and bronchitis are the most frequently recognized illness associated with *C. pneumoniae*, although asymptomatic infection or unrecognized mildly symptomatic illness are the most common result of infection. There are no specific symptoms and signs unique to pulmonary infections with this pathogen.<sup>(6)</sup> However, there are some clinical presentations associated with this infection. Although some cases have an acute onset, a more gradual onset is typical. Pharyngitis, sometimes with hoarseness is often present early in the course of the illness, may regress spontaneously

and cough with signs of lower respiratory tract infection develops from several days to weeks after the first symptoms giving the appearance of a biphasic pattern. Cough is often prolonged. There may be a history of fever, but fever is often not present at examination. Sinusitis, otitis media either as one of the initial complaints, or as a later complication may accompany *C. pneumoniae* infection.<sup>(16)</sup> If the lower respiratory tract is involved, rhonchi and rales are often heard on auscultation. A chest radiograph usually demonstrates a single subsegmental pneumonitis in milder cases, bilateral pneumonitis or pleural effusions have also been demonstrated in more severe cases.

Most *C. pneumoniae* associated respiratory illness are relatively mild both in children and adults. *C. pneumoniae* pneumonia usually responds to outpatient treatment and seldom requires hospitalization. Persons who are hospitalized are often older and have chronic disease. In some cases, however *C. pneumoniae* can cause severe illness, even in the absence of underlying disease. *C. pneumoniae* was isolated from the respiratory tract and pleural fluid of a previously healthy adolescent boy with severe pneumonia complicated by respiratory failure.<sup>(17)</sup>

Although the association of *C. pneumoniae* with pneumonia in our patient depends on serology, for the following reasons we believe that *C. pneumoniae* is causally related to the acute pneumonia syndrome in this child. *M. pneumoniae* infection which is the common cause of acute community acquired pneumonia is unlikely due to 1) the negative serological test in paired serum and a negative PCR test, 2) the high titer of IgM antibodies in either the acute or convalescent serum. IgM  $\geq 16$  has been proposed by Grayston et al. as the criterion for serological diagnosis of acute

*C. pneumoniae* infection.<sup>(1)</sup> In primary *C. pneumoniae* infection the *C. pneumoniae* MIF IgM antibody usually appears 3 weeks after the onset of illness. The IgG antibody titer may not appear until 6 to 8 weeks after onset. In case of reinfection, IgM antibody may not appear or may appear only at low titers, the IgG antibody titer rises quickly, often within 1 to 2 weeks, and may reach a value of 512 or above. 3) Although only a twofold rise in titer has been demonstrated, this may be due to premature detection of antibody response in the course of illness, and early treatment with specific antibiotics may have modified the antibody response.

It has been reported that symptoms associated with *C. pneumoniae* infection may be of long duration in adults and children<sup>(18,19)</sup> which is in contrast to our patient. If symptoms persist after one course of antibiotics, a repeated course is recommended.<sup>(1,2)</sup> Some reports have suggested that prolonged courses for up to 3 weeks may be required to eradicate *C. pneumoniae* from the nasopharynx of adults with flu-like illness and pharyngitis.<sup>(20)</sup> Erythromycin, tetracycline and doxycycline are recommended for treatment in adults where as erythromycin is suitable for children. The new macrolides, roxithromycin, azithromycin and clarithromycin have been demonstrated to be effective against *C. pneumoniae* in vitro.<sup>(21,22)</sup> There have been reports of roxithromycin in treating *C. pneumoniae* pneumonia in adults with good clinical results.<sup>(23, 24)</sup> One multicenter study compared erythromycin suspension 40 mg/kg/day with clarithromycin suspension 15 mg/kg/day, for 10 days, in children between 3 and 12 years of age with radiologically proven pneumonia. Both drugs were equally

efficacious, yielding a clinical success rate of 95 and 98 percent, and eradicating the organism in 86 and 79 percent of the children, respectively.<sup>(14)</sup>

In the out - patient clinic it is not practical to evaluate the role of viruses as copathogens which may have contributed to the respiratory illness of our patient with *C. pneumoniae* infection. Previous studies in developed countries.<sup>(25)</sup> have shown evidence of mixed etiology in acute lower respiratory tract infections of children

*C. pneumoniae* also may act as an inflammatory trigger for asthma. There have been reports of patients with culture-documented *C. pneumoniae* infection who developed significant bronchospasm.<sup>(19,26)</sup> Emre et al.<sup>(27)</sup> reported an association of *C. pneumoniae* infection and reactive airway disease in children. There is evidence suggesting that the bronchial reactivity observed with *C. pneumoniae* infection may be IgE- mediated.<sup>(28)</sup> The potential of *C. pneumoniae* to cause prolonged and persistent infection may produce chronic inflammation and trigger bronchospasm in susceptible persons. Therefore, early recognition and treatment may prevent chronic lung infection. We also suggest that *C. pneumoniae* may be an important agent to consider as the causative organism of respiratory tract infection in children.

#### Acknowledgement

We also would like to thank Mr. P. Hirsch for editing the manuscript.

#### References

1. Kuo CC, Jackson LA, Campbell LA, Grayston JT. Chlamydia pneumoniae (TWAR) Clin Microbiol Rev 1995 Oct; 8(4): 451-61

2. Emre U, Roblin PM, Gelling M, Dumornay W, Rao M, Hammerschlag MR, Schachter J. The association of Chlamydia pneumoniae infection and reactive airway disease in children. Arch Pediatr Adolesc Med 1994 Jul; 148(7): 727-32
3. Saikku P, Mattila K, Nieminen MS, Makela PH, Huttunen JK, Valtonen V. Serological evidence of an association of a novel chlamydia, TWAR with chronic coronary heart disease and acute myocardial infarction. Lancet 1988 Oct 29; 2(8618): 983-6
4. Linnanmaki E, Leinonen M, Mattila K, Nieminen MS, Valtonen V, Saikku P. Chlamydia pneumoniae-specific circulating immune complexes in patients with chronic coronary heart disease. Circulation 1993 Apr; 87(4): 1130-4
5. Puolakkainen M, Kuo CC, Shor KA, Wang SP, Grayston JT, Campbell LA. Serological response to Chlamydia pneumoniae in adults with coronary arterial fatty streaks and fibrolipid plaques. J Clin Microbiol 1993 Aug; 31(8): 2212-4
6. Grayston JT. Chlamydia pneumoniae (TWAR) infections in children. Pediatr Infect Dis J 1994 Aug; 13(8): 675-85
7. Nunthapisud P, Tatiyakavee K, Prapphal N, Likitnukul S, Grayston JT. The prevalence of Chlamydia pneumoniae, strain TWAR, antibody in school children: a preliminary study. Chula Med J 1992 Apr; 36(4): 273-7
8. Saikku P, Ruutu R, Leinonen M, Panelius J, Tupasi TE, Grayston JT. Acute lower-respiratory-tract infection associated with chlamydia TWAR antibody in Filipino children. J Infect Dis 1988 Nov; 158(5): 1095-7
9. Thom DH, Grayston JT, Wang SP, Kuo CC, Altman J. Chlamydia pneumoniae strain TWAR, Mycoplasma pneumoniae, and viral infections in acute respiratory disease in a university student health clinical population. Am J Epidemiol 1990 Aug; 132(2): 248-56
10. Chirgwin K, Roblin PM, Gelling M, Hammerschlag MR, Schachter J. Infection with Chlamydia pneumoniae in Brooklyn. J Infect Dis 1991 Apr; 163(4): 757-61
11. Forgie IM, O'Neill KP, Lloyd-Evans N, Leinonen M, Compbell H, Whittle HC, Greenwood BM. Etiology of acute lower respiratory tract infections in Gambian children. II. Acute lower respiratory tract infection tract infection in children ages one to nine years presenting at the hospital. J Pediatr Infect Dis 1991 Jan; 10(1): 42-7
12. Grayston JT, Aldous MB, Easton A, Wang SP, Kuo CC, Campbell LA, Altman J. Evidence that Chlamydia pneumoniae cause pneumonia and bronchitis. J Infect Dis 1993 Nov; 168(5): 1231-5
13. Gaydos CA, Eiden JJ, Oldach D, Mundy LM, Auwaerter P, Warner LM, Vance E, Burton AA, Quinn TC. Diagnosis of Chlamydia pneumoniae infection in patients with community-acquired pneumoniae by polymerase chain reaction enzyme immunoassay. Clin Infect Dis 1994 Jul; 19(1): 157-60
14. Block S, Hedrick J, Hammerschlag MR, Cassell GH, Craft JC. Mycoplasma pneumoniae and Chlamydia pneumoniae in pediatric community-acquired pneumonia: comparative efficacy

- and safety of clarithromycin vs erythromycin ethyl-succinate. *Pediatr Infect Dis J* 1995 Jun; 14(16):471 - 7
15. Jantos CA, Wienphal B, Schiefer HG, Wagner F, Hegemann JH. Infection with Chlamydia pneumoniae in infants and children with acute lower respiratory tract disease. *Pediatr Infect Dis J* 1995 Feb; 14(2): 117 - 22
16. Normann E, Gnarpe J, Gnarpe H, Wettergren B. Chlamydia pneumoniae in children with acute respiratory tract infection. *Acta Paediatr* 1998 Jan; 87(1): 23 - 7
17. Augenbraun MH, Roblin PM, Mandel LJ, Hammerschlag MR, Schachter J. Chlamydia pneumoniae pneumonia with pleural effusion: diagnosis by culture. *Am J Med* 1991 Oct; 91(4): 437 - 8
18. Falck G, Heyman L, Gnarpe J, Gnarpe H. Chlamydia pneumoniae and chronic pharyngitis. *Scand J Infect Dis* 1995 ; 27(2): 179 - 82
19. Normann E, Gnarpe J, Gnarpe H. Wettergren B. Chlamydia pneumoniae in children with acute respiratory tract infection. *Acta Paediatr* 1998 Jan; 87(1): 23 - 7
20. Hammerschlag MR, Chirgwin K, Roblin PM, Gelling M, Dumornay W, Mandel L, Smith P, Schachter J. Persistent infection with Chlamydia pneumoniae following acute respiratory illness. *Clin Infect Dis* 1992 Jan; 14(1): 178 - 82
21. Hammerschlag MR, Qumei KK, Roblin PM. In vitro activities of azithromycin, clarithromycin, ofloxacin, and other antibiotics against Chlamydia pneumoniae. *Antimicrob Agents Chemother* 1992 Jul; 36(7): 1573 - 4
22. Freidank HM, Losch P, Vogele H, Wiedmann-Al-Ahmad M. In vitro susceptibilities of Chlamydia pneumoniae isolated from German patients and synergistic activity of antibiotic combinations. *Antimicrob Agents Chemother* 1999 Jul; 43(7): 1808 - 10
23. Chikumi H, Kanamoto Y, Matsumoto Y, Sugimoto Y, Sakata T, Teramoto H, Yamasaki S, Konichi T, Tatsukawa T, Sasaki T. A case of adult Chlamydia pneumoniae pneumonia diagnosed from a culture method. *Kansenshogaku Zasshi J Japan Assoc Infect Dis* 1994 Mar; 68(3): 407 - 10
24. Ducroix JP, Smail A, Lafon B, Devendeville A, Orfila FFB, Baillet J. Treatment with roxithromycin of community-acquired pneumonia due to Chlamydia pneumoniae. Abstract presented at 19<sup>th</sup> international congress of chemotherapy July 16 - 21 1995 Montreal Quebec, Canada.
25. Paisley JW, Lauer BA, McIntosh K, Glode MP, Schachter J, Rumack C. Pathogens associated with acute lower respiratory tract infection in young children. *Pediatr Infect Dis* 1984 Jan-Feb; 3(1): 14 - 9
26. Hahn DL, Dodge RW, Golubjatnikov R. Association of Chlamydia pneumoniae (strain TWAR) infection with wheezing asthmatic bronchitis, and adult-onset asthma. *JAMA* 1991 Jul 10; 266(2): 225 - 30
27. Emre U, Roblin PM, Gelling M, Dumornay W, Rao M, Hammerschlag MR, Schachter J. The association of Chlamydia pneumoniae infection and reactive airway disease in children 1944. *Arch Pediatr Adolesc Med* 1994 Jul; 148(7): 727 - 32

28. Emre U, Sokolovskaya N, Roblin PM, Schachter J, Hammerschlag MR. Detection of anti-Chlamydia pneumoniae IgE in children with reactive airway disease. *J Infect Dis* 1995 Jul; 172(1): 265-7