

Original article

Reliability and validity of the family state and functioning assessment scale

Suchat Supphapitiphon^{a,*}, Napakkawat Buathong^b, Siriluck Suppapitiporn^c

^aDepartment of Family Medicine, King Chulalongkorn Memorial Hospital, Thai Red Cross Society, Bangkok, Thailand

^bDepartment of Family Medicine and Preventive Medicine, Faculty of Medicine, Prince of Songkla University, Songkhla, Thailand

^cDepartment of Psychiatry, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

Background: Family significantly influences health of family members in both positive and negative ways. There are limited instruments in Thailand for assessing family state and functioning on adult patients.

Objectives: To examine the validity and reliability of the family state and functioning assessment scale (FSFAS) which was designed to assess the extent one perceived about family issues and functioning.

Methods: The scale was conducted on 1,200 Thai adults: 800 outpatients attending the hospitals and 400 participants in community. Of the sample, 70% were women and the mean age was 50.4 years. The psychometric properties of the scale were examined in terms of construct validity and reliability. Exploratory factor analysis with principal components analysis and varimax rotation was performed to assess factor structures. Cronbach's alpha coefficient was calculated to estimate reliability.

Results: Exploratory factor analysis showed that the scale had a five-factor structure (support, discipline, communication and problem solving, emotional status and relationship) that accounted for 57.3% of the total variance. The final version of the scale consisted of 25 items with a Cronbach's alpha coefficient of 0.87 for the total and 0.70 - 0.84 for the subscales.

Conclusions: The scale has acceptable factorial validity and internal consistency reliability which can be useful instrument for assessing family state and functioning.

Keywords: Family state, functioning, reliability, validity.

Family significantly impacts on health of family members in both positive and negative ways.⁽¹⁾ Having a close-knit and supportive family provides emotional support, economic well-being, and increases overall health. When family life is characterized by stress and conflict, the health of family members tends to be negatively affected. Therefore, evaluating the family issues and family functioning is important to help understand and encourage families to perform their duties effectively. These can be conducted by several ways such as semi-structured interviews, observing the interactions in the family or between family members. But such methods require a lot of time and

may not be practical in some clinical settings and research. Ease of administration and cost efficiency make self-report instruments attractive for assessing psychological constructs in large-scale research. Questionnaires are often developed for a particular purpose.⁽²⁾ Family APGAR^(3 - 5) is a 5-question assessment tool used for rapid assessment of family function and dysfunction. It measures individual's level of satisfaction about family relationships. The five dimensions of family satisfaction are: adaptation, partnership, growth, affection and resolve. The internal consistency is quite high ($r = 0.59 - 0.80$). However, many researchers did not find the consistency of the score with the assessment of family functioning by the therapist and did not find support for using the measurement tool.^(6 - 7)

McMaster family assessment device (FAD)^(8 - 9) consists of a 60-item self-report question-naire that evaluates 6 dimensions of family functioning and

*Correspondence to: Suchat Supphapitiphon, Department of Psychiatry, Faculty of Medicine, chulalongkorn University, Bangkok 10330, Thailand.

Received: March 10, 2018,

Revised: May 20, 2018,

Accepted: August 10, 2018

overall general family functioning. The 6 dimensions of family functioning are as follows: problem solving, communication, roles, affective responsiveness, affective involvement, and behavior control. Internal consistency was estimated at 0.83 - 0.90 and intercorrelation was 0.4 - 0.6. Discriminant analysis found statistical difference in family functioning between schizophrenic families and normal families. Family functioning has been implicated in the onset of child and adult psychopathology. FAD was translated into multiple languages and has been extensively used in a variety of research contexts and clinical practices.^(10 - 13) Family assessment measure (FAM) was developed for family assessments according to seven key dimensions: task accomplishment, role performance, communication, affective expression, involvement, control, values and norms. It contained 92 items and provides measures of the family as a system and of the relationships.⁽¹⁴⁻¹⁵⁾ As for Thailand, there are some instruments that measure family functioning and family health promoting behaviors such as the Chulalongkorn Family Inventory⁽¹⁶⁾, the Thai Family Functioning Scale⁽¹⁷⁾, the Thai Family Health Routines (TFHR) scale⁽¹⁸⁾, the Family Health Promoting Behavior Scale⁽¹⁹⁾ and the Perceived Family support.^(20 - 21) Chulalongkorn Family Inventory (CFI)⁽¹⁶⁾ by Trangkasombat U consisted of 36 items to study family functioning in the families of psychiatric patients compared with nonclinical families⁽²²⁾ and to examine differences in family functioning between the families of patients with depressive disorders and with schizophrenia.⁽²³⁾ Thai family functioning scale⁽¹⁷⁾ was reported to have better psychometric properties than the Thai version of the FAD; however, it was tested only on adolescents aged 15 to 19. Thai family health routines (TFHR) scale⁽¹⁸⁾ which used to measure the health of Thai families through their routine behaviors in daily life comprises 70 items. No instrument is now available for quick, simple evaluating family issues and functioning in Thai medical outpatients and community. Developing such an instrument is necessary for assessing family functioning and screening to identify families experiencing problems, in view of the increasing role of families in care for medical patients. Therefore, this study was aimed to examine the validity and reliability of the family state and functioning assessment scale (FSFAS) which was designed to assess to what extent one perceived the family issues and functioning.

Methods

Scale development and initial reliability estimates

A literature search and review was conducted covering the area of interest, family functioning assessment in medical patients.^(2 - 21) Development of the family state and functioning assessment scale (FSFAS) was mainly based on the structural domains of Family Assessment Device from McMaster Model of Family Functioning (MMFF)⁽⁸⁾, Chulalongkorn family inventory⁽¹⁶⁾ and the Thai family functioning scale.⁽¹⁷⁾ The questionnaire was used to answer issues and functioning about emotional, support, communication and social interaction among members of the household. The scale was validated by content experts (family physicians, psychiatrists, psychologists and nurses) and initially composed of 30 items. Preliminary pilot testing was conducted on 30 participants. The Cronbach's alpha coefficient was 0.7. Five statements were adjusted for better understanding.

Scale validation and studying in participants

A cross-sectional descriptive study was conducted. The sample size must meet an optimal sample size for factor analysis that required respondents at least ten times the number of items⁽²⁴⁾ and a sample size of at least 1,000 might be appropriate for exploratory factor analysis.⁽²⁵⁾ For reducing the error we increased the sample size with 20% so the scale was conducted on 1,200 Thai adults aged over 18 years. Due to the sample should be chosen to be as similar as possible to the relevant population, thus eight hundred outpatients from 2 hospitals and 400 participants from 2 communities in Bangkok, Thailand were asked to participate in the study. Data were collected using self-administered demographics questionnaire and the family state and functioning assessment scale (FSFAS).

FSFAS is a self-reporting questionnaire asking to indicate the degree to which they agree with various statements on a four-point Likert scale ranging from "1 = strongly disagree" to "4 = strongly agree". The scale is composed of both positive and negative statements, and recoding score on negative statement items should be done before calculating the total score. The positive statements are items number 2, 4, 5, 6, 8, 9, 10, 12, 17, 20, 23, 24, 25, 26, 27, 29, and 30; and, the negative statements are items number 1, 3, 7, 11, 13, 14, 15, 16, 18, 19, 21, 22, and 28. The total score of the FSFAS is obtained by summing raw scores across 30

items that can range from 30 to 120 and calculating the mean score. A higher score indicates a better family state and functioning.

Statistical analysis

In order to summarize the characteristics of the participants, descriptive statistics, percentage and frequency were used for categorical variables and mean and standard deviation were calculated for continuous variables. The internal consistency reliability was assessed using Cronbach’s alpha. The construct validity was analyzed using exploratory factor analysis which was conducted by principal components extraction, followed by varimax rotation. Eigenvalues, relative magnitude and direction of factor loadings explaining variance and communality, were examined in these analyses. Kaiser’s Eigen value greater than 1 was used to determine the number of factors. Significance was set at alpha = 0.05; two-tailed for all statistical tests.

Results

The study population consisted of adults aged 18 – 90 years, with a mean age of 50.4 years (SD 16.1). Approximately 70% were women. Nearly half were married or living as married couples and 31% were single. About 29% had a primary school, 24% had a high school diploma and 28.3% had a bachelor degree. Data are shown in Table 1.

The mean score of each item of the FSFAS ranged from 2.20 to 3.38(SD 0.80 - 1.02). The mean score of 30-item FSFAS was 84.32 (SD 11.13). The internal consistency analysis for the scale had a Cronbach’s alpha coefficient of 0.82. Four items (item number 3, 18, 19 and 22) had corrected item-total correlation less than 0.2. After removing these items by proceeding 1 item at a time, and redone the reliability analysis after each deleted item, the scale had higher Cronbach’s alpha coefficient of 0.84 - 0.86 as presented in Table 2.

Table 1. Demographic data of participants.

Variables	Number	Percentage
Sex (n = 1,174)		
Male	346	29.5
Female	828	70.5
Age (n = 1,158)		
≤ 25 years	106	9.1
26 - 35	132	11.4
36 - 45	178	15.4
46 - 55	265	22.9
> 55	477	41.2
Mean (SD) = 50.4 years(16.1), range = 18 - 90		
Marital status (n = 1,185)		
Married	564	47.6
Single	369	31.1
Divorced/separated	161	21.3
Educational level (n = 1,182)		
No school	19	1.6
Primary school	344	29.1
Secondary school	144	12.2
High school or diploma	284	24.1
Bachelor degree	335	28.3
Master degree and above	56	4.7
Occupation (n = 1,176)		
Unemployed	264	22.5
Employee	323	27.5
Business	278	23.6
Government officer	147	12.5
Others	164	13.9

Table 2. Reliability analysis based on the corrected item-total correlation (CITC) and Cronbach’s alpha coefficient if item was deleted.

Item	Mean	S.D.	CITC	Cronbach’s Alpha if Item Deleted
3.	2.53	0.83	0.042	0.837
18.	2.20	0.91	-0.564	0.857
19.	3.38	0.82	0.010	0.838
22.	2.60	0.96	0.066	0.838

Exploratory factor analysis

The structure of the FSFAS was analyzed using principal components extraction. As for the 26-item sets, the requirements for exploratory factor analysis in this sample were fulfilled (Kaiser-Meyer-Olkin measure of adequacy = .913, Bartlett’s test of sphericity $P < .001$). Communality of 26 items ranged 0.36 to 0.73. Using the initial factor solutions, items were removed step-by-step based on the following criteria: factor loading $< .40$, cross-loading $> .30$, communality $< .30$ and corrected item-scale correlation $< .30$.⁽²⁶⁾ According to these criteria five factors were extracted and the scale had 25 items with factor loading between 0.52 to 0.80. Each factor had eigenvalues greater than 1. Exploratory factor analysis showed that the scale had a five-factor structure that accounted for 57.3% of the total variance. Factor 1 was accounted for 13.8% of the variances, consisted of 5 items that we called family support (being able to access family members that a person can rely upon if needed). Factor 2 was accounted for 12.5% of the variances, consisted of 6 items that we called family discipline (the process to help family member learn appropriate behaviors and share responsibilities). Factor 3 was accounted for

11.8% of the variances, consisted of 5 items that we called communication and problem solving (how family members exchange verbal information and the family’s ability to resolve problems). Factor 4 was accounted for 10.6% of the variances, consisted of 5 items that we called emotional status (emotions that are shared with and between family members which measures the satisfaction with the intimacy and emotional interaction that exist in the family). Finally, the fifth factor was accounted for 8.5% of the variances, consisted of 4 items that we called family relationship (a person’s perception of the quality of his or her family relationship functioning). The total score of the 25-item FSFAS is scored by adding the responses (1- 4) for each scale and divided by the number of items in each scale (4 - 6). A higher score indicates a better family state and functioning (Table 3).

Reliability

The internal consistency analysis for total scale and each subscale were calculated. The final version of the scale consisted of 25 items with a Cronbach’s alpha coefficient of 0.87 for the total and 0.70 - 0.84 for the subscales (Table 4).

Table 3. Factor loadings from principal axis factor analysis with varimax rotation.

Component	Extraction Sums of Square Loading			Rotation Sums of Square Loading		
	Total (Eigenvalue)	% of Variance	Cumulative%	Total (Eigenvalue)	% of Variance	Cumulative%
1	7.476	28.755	28.755	3.589	13.802	13.802
2	3.925	15.098	43.853	3.247	12.487	26.289
3	1.344	5.168	49.020	3.077	11.833	38.122
4	1.133	4.357	53.378	2.765	10.636	48.759
5	1.021	3.927	57.304	2.222	8.546	57.304

Kaiser-Meyer-Olkin Measure of Sampling Adequacy = 0.913, Bartlett’s test of Sphericity ; $P < 0.001$

Table 4. Domains and Cronbach's alpha coefficient of the FSFAS-25 items.

Domains	Number of Items	Cronbach's alpha coefficient
Family support	5	0.84
Discipline	6	0.83
Communication and problem solving	5	0.82
Emotional status	5	0.70
Relationship	4	0.77
Total	25	0.87

family support : being able to access family members that a person can rely upon if needed; family discipline : the process to help family member learn appropriate behaviors and share responsibilities communication and problem solving : how family members exchange verbal information and the family's ability to resolve problems ; emotional status : emotions that are shared with and between family members which measures the satisfaction with the intimacy and emotional interaction that exist in the family ; family relationship : a person's perception of the quality of his or her family relationship functioning.

Discussion

The study was carried out to examine the psychometric properties of the family state and functioning assessment scale, in a sample of Thai adults from community and hospital settings. The results indicate that the FSFAS is a psychometrically sound instrument.

According to exploratory factor analysis, Kaiser-Meyer-Olkin Measure of Sampling Adequacy – this measure varies between 0 and 1, and values closer to 1 are better. A value of .6 is a suggested minimum. This scale had KOM = .913 and significant Bartlett's test of sphericity ($P < .001$). These tests showed good results to conduct principal components analysis. Principal components analysis is a method of data reduction that requires a large sample size. This scale was conducted on 1,200 Thai adults that met an optimal sample size for factor analysis since at least 1,000 families were needed for the instrument evaluation. The structure of the FSFAS was analyzed using principal components extraction. The results showed that the scale had a five-factor structure that accounted for 57.3% of the total variance. It consisted of 25 items with factor loading between 0.52 to 0.80. Each factor had eigenvalue greater than 1. These indicated that the scale had acceptable factorial validity.⁽²⁶⁾

The final version of the scale consisted of 25 items with a Cronbach's alpha coefficient of 0.87 for the total and 0.70 - 0.84 for the subscales. A high value of total alpha indicates good reliability. The internal consistency of this total scale was closely similar to the previous studies.^(8 - 9, 16)

This short scale which measures family functioning is appropriate for use in clinical practice and research. It can assist the clinician in determining how to build on current family strengths, as well as identify areas for growth that could be beneficial for promoting family functioning effectively. Family assessment can be used for early identification of patients at risk for poor family functioning and screening to identify families experiencing problems. Furthermore, it may be possible to improve medical condition and adherence by working with the family in specific areas of family functioning.^(27 - 29) Nevertheless, this study had some limitations that should be noted. First, the measures we used were limited as they relied on participants' self-reports. Second, the proportion of women was higher than that of men in this study. It is possible that women have slightly outnumbered the men in the total population for Thailand and might be more willing to participate in the study. However, we enrolled more than three hundred men that was also an adequate sample size for the representativeness of the relevant population. Finally, the sample consisted of participants mostly living in Bangkok. Further studies should be conducted in populations from other regions of Thailand.

Conclusions

FSFAS has acceptable factorial validity and internal consistency reliability that can be used as a quick and effective tool in both clinical practice and research to assess the extent one perceived about the family issues and functioning. These would provide valuable information for promoting effective

functioning of families and hence improving physical and mental health status.

Acknowledgements

This study was part of the research project 'depression, quality of life and model of service consistent with the needs of community and care setting' funded by the Ratchadapisek Sompotch Fund of Chulalongkorn University, Thailand. The authors would like to our thanks to the hospital staff and healthcare personnel for their assistance with data collection, content experts for scale development and all respondents for their participations in the study.

Conflict of interest

None of the authors has any potential conflict of interest to disclose.

References

- Ross C, Mirowsky, J, Goldstein K. The impact of the family on health: The decade in review. *J Marriage Fam* 1990;52:1059-78.
- Pritchett R, Kemp J, Wilson P, Minnis H, Bryce G, Gillberg C. Quick, simple measures of family relationships for use in clinical practice and research. A systematic review. *Fam Pract* 2011;28:172-87.
- Smilkstein G. The family APGAR: a proposal for a family function test and its use by physicians. *J Fam Pract* 1978;6:1231-9.
- Good MJ, Smilkstein G, Good BJ, Shaffer T, Arrons T. The family APGAR index: a study of construct validity. *J Family Pract* 1979;8:577-82.
- Smilkstein G, Ashworth C, Montano D. Validity and reliability of the family APGAR as a test of family function. *J Fam Pract* 1982;15:303-11.
- Mengel M. The use of the family APGAR in screening for family dysfunction in a family practice center. *J Fam Pract* 1987;24:394-8.
- Gardner W, Nutting PA, Kelleher KJ, Werner JJ, Farley T, Stewart L, et al. Does the family APGAR effectively measure family functioning? *J Fam Pract* 2001;50:19-25.
- Epstein NB, Bishop DS, Leven S. The McMaster model of family functioning. *J Marriage Fam Couns* 1978;4:19-31.
- Epstein NB, Baldwin LM, Bishop DS. The McMaster family assessment device. *J Marital Fam Ther* 1983;9:171-80.
- Staccini L, Tomba E, Grandi S, Keitner GI. The evaluation of family functioning by the family assessment device: a systematic review of studies in adult clinical populations. *Fam Process* 2015;54:94-115.
- Mansfield AK, Keitner GI, Dealy J. The family assessment device: an update. *Fam Process* 2015;54:82-93.
- Hamilton E, Carr A. Systematic review of self-report family assessment measures. *Fam Process* 2016;55:16-30.
- Botelho de Haan KL, Hafekost J, Lawrence D, Sawyer MG, Zubrick SR. Reliability and validity of a short version of the general functioning subscale of the McMaster Family Assessment Device. *Fam Process* 2015;54:116-23.
- Skinner H, Steinhauer P, Santa-Barbara J. The family assessment measure. *Can J Community Ment Health* 1983;2:91-105.
- Skinner H, Steinhauer P, Sitarenios G. Family Assessment Measure (FAM) and process model of family functioning. *J Fam Ther* 2000;22:190-10.
- Trangkasombat U. Family functioning. In: Trangkasombat U, ed. *Family therapy and family counseling*. 1sted. Bangkok: Fuangfa Printing, 1997: 38-54.
- Suttiamnuaykul W. Measuring family functioning in Thailand: Developing the Thai Family Functioning Scale TFFS and comparing its psychometric properties to those of the Thai version of the Family Assessment Device FAD. *Siriraj Hospital Gazette* 2003;55(Suppl 1): 75.
- Kanjanawetang J, Yunibhand J, Chaiyawat W, Wu YW, Denham SA. Thai Family Health Routines: scale development and psychometric testing. *Southeast Asian J Trop Med Public Health* 2009;40: 629-43.
- Suwanpatikorn K. Family Health Promoting Behavior Scale: development and psychometric analysis [Dissertation], Bangkok: Mahidol University; 2001 [cited 2015 July 1]. Available from: <http://www.thaithesis.org/detail.php?id=45458>.
- Anongrsuksai N. Relationship between family function and perceived family support with diabetes self-care behavior of patients with type 2 diabetes mellitus in Banpong District, Ratchaburi Province. *Nursing Journal of the Ministry of Public Health* 2013; 23:60-9.
- Zhang H. Family support and self-care behavior of breast cancer patients receiving combined therapy at six teaching hospital Meijing. Chiang Mai: Chang Mai University; 1999.
- Trangkasombat U. Family functioning in the families

- of psychiatric patients: a comparison with nonclinical families. *J Med Assoc Thai* 2006;89:1946-53.
23. Trangkasombat U. Family Functioning in Mental Illness: A Study in Thai Families with Depressive Disorders and Schizophrenia. *J Fam Psychother* 2008;19:1-23.
 24. Comrey AL, Lee HB. *A first course in factor analysis*. Hillsdale, NJ: Erlbaum: 1992.
 25. Pett MA, Lackey NR, Sullivan JJ. *Making sense of factor analysis: The use of factor analysis for instrument development in health care research*. California: Sage Publications; 2003.
 26. Worthington RL, Whittaker TA. Scale development research: a content analysis and recommendations for best practice. *Counsel Psychol* 2006;34:806-38.
 27. Garcia-Huidobro D, Puschel K, Soto G. Family functioning style and health: opportunities for health prevention in primary care. *Br J Gen Pract* 2012;62:e198-e203.
 28. Miller TA, Dimatteo MR. Importance of family/social support and impact on adherence to diabetic therapy. *Diabetes Metab Syndr Obes* 2013;6:421-6.
 29. Hayaki C, Anno K, Shibata M, Iwaki R, Kawata H, Sudo N, et al. Family dysfunction: A comparison of chronic widespread pain and chronic localized pain. *Medicine (Baltimore)* 2016;95:e5495.