

## Original article

# Depression in patients with type 2 diabetes attending at Chao Phya Abhaibhubejhr Hospital

Amporn Nueangjaknak<sup>a</sup>, Siriluck Suppakitiporn<sup>b,\*</sup><sup>a</sup> Program in Mental Health, Department of Psychiatry, Faculty of Medicine, Chulalongkorn University<sup>b</sup> Department of Psychiatry, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

**Background:** Diabetes mellitus (DM) is one of the most common public health problems worldwide. It is a chronic, debilitating disease that needs a lifelong care for both physical and psychological aspects. Previous studies on depression in diabetic patients are scant in Thailand.

**Objective:** To determine the prevalence of depression and its associated factors in patients with type 2 diabetes at Chao Phya Abhaibhubejhr Hospital, Prachinburi Province.

**Methods:** A cross-sectional descriptive study was conducted among patients with type 2 diabetes at the Outpatient Clinic of Chao Phya Abhaibhubejhr Hospital, Prachinburi Province, Thailand from August to December 2017. The instruments were demographic and clinical characteristics questionnaires, the Thai Hospital Anxiety and Depression Scale (Thai HADS), Health Perception Questionnaire, and family status and functioning assessment scale. Statistics used to analyze data were frequencies, percentages, mean and standard deviation, Chi-square test, Fisher exact test, Pearson product-moment correlation coefficient and logistic regression.

**Results:** Of the total 355 participants, the mean age was 55.7 years, 60 % had been diagnosed type 2 diabetes for more than 5 years; 25 % had used traditional herbs. The prevalence of depression in patients with type 2 diabetes was 20.8 %. Depression was significantly related to the age group above 50 years old, unemployment, licorice herb use and lower family functioning.

**Conclusion:** One-fifth of the patients with type 2 diabetes were found to have depression. The results of this study would be beneficial for recognizing depression and providing appropriate care for patients with type 2 diabetes.

**Keywords:** Depression, diabetes type 2, licorice, family functioning.

Diabetes mellitus type 2 is one of the most common chronic diseases;<sup>(1)</sup> it imposes an increasing economic burden on national healthcare systems worldwide.<sup>(2)</sup> The data represented by the National Survey of Risk Factors for Non-Communicable Diseases of Thailand indicate that the number of patients suffering from diabetes is 8.9%, equal to 4 million patients. The disease is associated with serious complications and commonly found comorbidities which affect one's health.<sup>(3)</sup> The disease also has a great impact on psychological and social performance of diabetic patients. Depression is a common mental

disorder characterized by sadness, loss of activities and by decreased energy. It is more common in medically ill individuals including diabetes. Patients with type 2 diabetes mellitus often have depression<sup>(4)</sup>, impaired family functioning and quality of life.<sup>(5,6)</sup> The presence of diabetes doubles the odds of comorbid depression. The prevalence of comorbid depression was significantly higher in diabetic women (28%) than in diabetic men (18%).<sup>(4)</sup> Wang J, *et al.* reported a significantly negative relationship between the family function and quality of life in Chinese patients with diabetes type 2.<sup>(5)</sup> Significant evidence supports a range of interventions and strategies to improve diabetes outcome.<sup>(7)</sup> Thailand is facing increasing numbers of patients with diabetes and the Thai government recognizes the burden of diabetes and has launched nationwide programs of health promotion and disease prevention.<sup>(8)</sup> Addressing the psychological

\*Correspondence to: Siriluck Suppakitiporn, Department of Psychiatry, Faculty of Medicine, Chulalongkorn University, Bangkok 10330, Thailand.

Received: June 18, 2018

Revised: August 31, 2018

Accepted: November 6, 2018

and family factors related to diabetes is very important. Understanding depression would be expected to produce substantial health benefits and improve outcomes of patients with diabetes in Thailand. Prachinburi Province was found to have the second highest prevalence rate of diabetes in Thailand. Therefore, this study was conducted to determine the prevalence of depression and associated factors in patients with type 2 diabetes at Chao Phya Abhaibhubejhr Hospital, Prachinburi Province.

## Methods

A cross-sectional study was conducted at Outpatient Clinic of Chao Phya Abhaibhubejhr Hospital, Prachinburi Province, Thailand. The recruited subjects were 355 adults with type 2 diabetes attending the Outpatient Clinic, Chao Phya Abhaibhubejhr Hospital from August to December 2017. They were able to understand Thai language and willing to participate in the study. Systematic random sampling was employed by randomly choosing a number to start with. The patients who received the queue numbered would be prioritized for selection; then every four persons from would be recruited until the sample size was reached. The study has been approved by the Institutional Review Board (IRB) of the Faculty of Medicine, Chulalongkorn University IRB NO.179/60 and of that of the Chao Phya Abhaibhubejhr Hospital. A written informed consent was obtained from each subject who agreed to participate in the study. Self-administered questionnaires were used and fasting blood glucose level was recorded.

## Measures

Demographic information (such as age, gender, marital status, educational level, occupation), clinical characteristics (regarding medical illnesses, duration of diabetes, history of substances and herbal use) and other relevant information were assessed. Depression was assessed by Hospital Anxiety and Depression Scale (HADS). The HADS was developed by Zigmond S, *et al.*<sup>(9)</sup> and the Thai version of HADS was developed by Nilchaikovit T, *et al.*<sup>(10)</sup> It comprises fourteen items for two sub-scales. Each of the two sub-scales relates to anxiety and depression and consists of seven items which obtain responses on a four-point Likert scale, ranging from 0 = mostly to 3 = not at all. Possible sub-scale scores range from 0 to 21. Even numbered questions are for depression. Recommended cut-off scores according to Zigmond S, *et al.*<sup>(8)</sup> are 8 – 10 for doubtful cases and 11 or

above for definite cases. An optimal balance between sensitivity and specificity was found using a cut-off score of 8 or above for HADS Depression.<sup>(11)</sup> For statistical purposes, HADS scores on depression subscale were calculated and participants with a depression score 8 or more were considered depression.<sup>(10)</sup> The Cronbach's alpha coefficient of depression sub-scale was 0.83.

Health perception was assessed by Health Perception Questionnaire.<sup>(12)</sup> This scale was developed by Chaninphon Temrat for her research. It was modified from the health belief questionnaire of Wongwairoj A.<sup>(13)</sup> that based on the concept of health belief model by Becker MH, *et al.*<sup>(14)</sup> The questionnaire consisted of 20 questions related to 4 aspects of health perception. Moreover, 8 questions related to diabetes perception were also added to the questionnaire. Therefore, this questionnaire contained 28 questions in total. The content validity was verified by 3 specialists: a psychiatrist, a general practitioner and a nurse; and the Cronbach's Alpha coefficient of the research tool was 0.84. Each question of the questionnaire was on a 5-point Likert response from 1 = strongly disagree to 5 = strongly agree. The results from sum of the total items range from 5 to 140 and were categorized into 3 levels of health perception: 50% and lower indicate mild; 51% - 69% indicate moderate; and 70% - 100% indicate high.

Family functioning was assessed by family state and functioning assessment scale (FSFAS)<sup>(15)</sup> that was based on Family Assessment Device (FAD) questionnaire designed by Epstein NB, *et al.*<sup>(16, 17)</sup> and Chulalongkorn Family Inventory by Trangkasombat U.<sup>(18)</sup> This instrument appraises five family factors on a 4 - Likert response from "1 = strongly disagree" to "4 = fully agree". The scale is composed of both positive and negative statements, and recoding score on negative statement items should be done before calculating the total score. The positive statements are items number 2, 4, 5, 6, 8, 9, 10, 12, 17, 20, 23, 24, 25, 26, 27, 29, 30 and the negative statements are the rest items. The total score of the FSFAS is obtained by summing raw scores across 30 items that can range from 30 to 120 and then can also calculate the mean score ranged from 1 to 4. The higher the score indicates the better the family functioning. Cronbach's alpha coefficient of this test in the Epstein NB, *et al.* study was 0.92, in the Trangkasombat study was 0.8, in the Suppapatiporn study was 0.87 and in the present study was 0.8.

**Statistical analyses**

To summarize the characteristics of the participants, descriptive statistics, percentage and frequency were used for categorical variables and mean, standard deviation were calculated for continuous variables. Chi-square test, Fisher exact test, Pearson product-moment correlation coefficient and logistic regression were used to explore the relationship between depression and other variables. All statistical tests were considered significant at *P* - value < 0.05.

**Results**

Demographic and clinical characteristics of the study sample are shown in Table 1. The mean age of

the participants was 55.7 years (SD = 13.6). The majority of the patients (78.9%) had medical comorbidities such as hypertension, dyslipidemia and renal problem. Herbal use was found in 25% of the sample. In these, 43 participants used curcuma (12.1%) and 10 used licorice herb (2.8%). More than 60% had fasting blood glucose level above 126 mg/dl. Of the sample (355), 52 participants had the HADS score for depression between 8 to10 and 22 participants had score of 11 or above so the prevalence of depression in patients with diabetes type 2 was 20.8 %. Most of them had moderate level of health perception (84.8%) and good family functioning (75.8%) (Table 2.).

**Table 1.** Socio-demographic and clinical characteristics of the participants. (n = 355)

Variables	Number	Percentage	
<b>Gender</b>	Male/ Female	163/192	45.9/54.1
<b>Age (years)</b> [mean 55.7 (SD13.6), min = 18, max = 89]			
	40 years or lower	48	13.5
	41 – 50	87	24.5
	51 - 60	95	26.8
	61-70	72	20.3
	above 70	53	14.9
<b>Educational</b>			
	No formal education	34	9.6
	Primary school	152	42.8
	Secondary school	56	15.8
	College or graduate or diploma	113	31.8
<b>Employment</b>			
	Unemployed	111	31.3
	Business	122	34.3
	Employee	39	11.0
	Agriculturist	29	8.2
	Government officer	29	8.2
	Others	25	7.0
<b>Comorbidities level</b>			
	No/ Yes	75/280	21.1/78.9
	Hypertension	229	64.5
	Dyslipidemia	203	57.2
	Renal disease	33	9.7
	Others	35	9.9
<b>Duration of diabetes (years)</b> [mean 9.2 (SD 7.6), min = 0.8, max = 50]			
	Less than 1	24	6.8
	1- 5	109	31.0
	5.1 - 10	118	33.5
	More than 10	101	28.7
<b>Blood glucose level (mg/dl)</b> [mean 153 (SD 51.9), min = 53, max = 400]			
	Less than 126	114	32.1
	126 or above	241	67.9
<b>Herbal use</b>			
	No/Yes	265/90	76.4/24.6
	Curcuma	43	12.1
	Bitter gourd	40	11.3
	Indian Gooseberry	34	9.6
	Licorice	10	2.8
	Others	5	4.2

**Table 2.** Frequency of participants by selected variables.

Variables	Number	Percentage
Thai-HADS [mean 4.6 (SD3.4), min = 0, max = 13]		
No depression (score 0 - 7)	281	79.2
Depression (score 8 or above)	74	20.8
Health perception [mean 73.7 (SD8.6), min = 40, max = 95]		
Mild (score < 50)	33	9.3
Moderate (51 - 69)	301	84.8
High (70 - 100)	21	5.9
Family functioning [mean 2.95 (SD 0.43), min = 2, max = 4]		
Fair (score 1.00 - 1.49)	0	0
Slightly good (1.50 - 2.49)	50	14.1
Good (2.50 - 3.49)	269	75.8
Very good (3.50 - 4.00)	36	10.1

Depression was significantly related to the age above 50 years, unemployment, licorice herb use and lower family state and functioning, as shown in Table 3. There was a significant correlation between depression and age, duration of diabetes, family functioning on overall, support, communication and problem solving, emotional state, and relationship (Table 4).

Further analysis using logistic regression showed that licorice, as a herb, unemployment, and family functioning (slightly good / good or very good) were independently associated with depression ( $P < 0.05$ ) (Table 5). Statistical analysis indicated no significant association between depression and fasting blood glucose level and other herbal use.

**Table 3.** Association between depression and variables (n = 355).

Variables	No depression	depression	X <sup>2</sup>	P - value
	N (%)	N (%)		
<b>Gender</b>				
Male	128 (78.5)	35 (21.5)	0.072	0.789
Female	153 (79.7)	39 (20.3)		
<b>Age (years)</b>				
≤ 50	115 (85.2)	20 (14.8)	4.801	0.028*
> 50	166 (75.5)	54 (24.5)		
<b>Employment</b>				
Unemployed	80 (72.1)	31 (27.9)	4.910	0.027*
Employed	201 (82.4)	43 (17.6)		
<b>Licorice use</b>				
Yes	5 (50)	5 (50)	5.301	0.036*
No	276 (80)	69 (20)		
<b>Family functioning</b>				
Slightly good	34 (68)	16 (32)	10.912	0.004**
Good	212 (78.8)	57 (21.1)		
Very good	35 (97.2)	1 (2.8)		

\* $P < 0.05$  \*\* $P < 0.01$

**Table 4.** Pearson correlation between depression score and variables.

Variables	<i>r</i>	<i>P</i> - value
Age (years)	0.199	<0.001**
Blood glucose level	0.023	0.668
Duration of diabetes	0.136	0.01*
Health perception	-0.080	1.310
Family functioning	-0.266	<0.001**
Family support	-0.194	<0.001**
Discipline	0.008	0.120
Communication and problem solving	-0.165	0.002**
Emotional state	-0.188	<0.001**
Relationship	-0.197	<0.001**

\**P* < 0.05 \*\**P* < 0.01

**Table 5.** Logistic regression for variables associated with depression.

Variables	$\beta$	S.E. ( $\beta$ )	<i>P</i> - value	Odds ratio (95%CI of OR)
Unemployment	0.59	0.27	0.028*	1.81 (1.06 - 3.07)
Licorice use	1.59	0.67	0.018*	4.91 (1.31-18.40)
Lower family functioning	0.69	0.33	0.039*	2.00 (1.03 - 3.87)
Constant	-1.542	0.131	<0.001**	

\**P* < 0.05 \*\**P* < 0.01

## Discussion

The study revealed that the prevalence of depression in patients with type 2 diabetes was 20.8 %. Depression was significantly related to age above 50 years, unemployment, licorice herb use and lower family functioning. The prevalence of depression in our study is comparable to studies using the same instrument by Engum A. (19%)<sup>(19)</sup>, Thaneerat T. (28%)<sup>(20)</sup> and Collin MM. (32%)<sup>(21)</sup>. The higher prevalence of depression was reported from studies among hospitalized diabetes patients (40 - 53.8%)<sup>(22, 23)</sup> while lower prevalence rate was found in primary care setting (11.5%)<sup>(24)</sup>. These have also demonstrated that depression is more common in diabetes patients compared with the general population.<sup>(25)</sup> Recognition of depression would be beneficial for providing appropriate care for patients with type 2 diabetes.

Age and unemployment were associated with depression. These findings are consistent with other studies which also found older age increase risk of depression<sup>(26, 27)</sup> and unemployment as well as retirement significantly associated with

depression.<sup>(28 - 30)</sup> It is likely that unemployed person lack feeling of stability<sup>(31)</sup>, lower self-esteem, and prone to have stressor about financial problem.

Our findings showed that there was a significant correlation between depression and family functioning. Family support, care and relationship are important parts of social support. Perceived support can be understood as a personal subjective evaluation that his or her social network will provide effective help when needed.<sup>(32)</sup> Due to diabetes is a chronic illness and usually accompanied by many comorbidities, thus assessment of family functioning and emphasizing the importance of appropriate family intervention to improve the family's competence in problem solving and to promote better communication in the family should be also addressed in diabetes patients for reducing depression. In addition, previous studies found that family and social support were important aspects of adherence to diabetes management<sup>(33)</sup> and good family function associated with better quality of life.<sup>(34)</sup> Moreover, some studies revealed that family functioning also related to control of diabetes<sup>(35)</sup> but we did not find the association of blood glucose level

with family functioning and depression. Therefore, individual circumstances, such as age, occupation, duration of diabetes and family factors must be considered and may lead to treatment targets and strategies to improve depression in diabetic patients.

This study showed that approximately 25% of the subjects had used herbs. Sawatsri P. revealed that nearly 60% of type 2 diabetic patients used herbs and indigenous food crop for blood glucose.<sup>(36)</sup> Herbal use was not associated with glycemic control in our study. However, the finding that licorice use was associated with depression and the likelihood of diabetes patients having depression was 4.9 times more in those with licorice use was of interest. A study on the effect of licorice on depression of nurses<sup>(37)</sup>, it seems that this drug can be used to reduce the depressive symptoms. The mechanism may be explained by the certain flavonoids extracted from licorice root inhibit the re-uptake of serotonin.<sup>(38)</sup> Limited data were available regarding licorice use in diabetes patients and its impact on depression. Therefore, further research is needed to explore the relationships and to show the reasons for our findings. In addition, providing correct information of herbal use for diabetes patients should be concerned with appropriate care for patients with type 2 diabetes.

Nevertheless, there are some limitations which need to be considered in this study. First, the study was conducted on a sample group of type 2 diabetes patients who attended the Diabetes Clinic of the Outpatient Department (OPD), Chao Phya Abhaibhubejhr Hospital, Prachinburi Province. As this hospital is a tertiary referral hospital, it may limit the extent to which these findings could be generalized to other hospitals with different settings. Second, since this study was a cross-sectional descriptive design, we could not identify the causal relationship between these variables. Third, the Thai Hospital Anxiety and Depression Scale is a screening tool, not usually used for diagnosis of mental disorder. Lastly, fasting blood glucose level was used for assessing glycemic control due to the limited data of blood HbA1c level.

## Conclusion

One-fifth of the patients with type 2 diabetes were found to have depression. Older age, unemployment, licorice herb use and lower family functioning were associated with depression. The results of this study would be beneficial for recognizing depression and providing appropriate care for patients with type 2 diabetes.

## Acknowledgments

The authors would like to express the gratitude to all the participants who were involved in this study and the staff of Chao Phya Abhaibhubejhr Hospital, Prachinburi Province, Thailand.

## Conflict of interest

None of the authors has any potential conflict of interest to disclose.

## References

1. National Diabetes Audit (NDA). The health and social care information report complication and mortality. Leeds, UK: NDA; 2012.
2. Zhang P, Zhang X, Brown J, Vistisen D, Sicree R, Shaw J, et al. Global healthcare expenditure on diabetes for 2010 and 2030. *Diabetes Res Clin Pract* 2010;87: 293-301.
3. Diabetes Association of Thailand under The Patronage of Her Royal Highness Princess Maha Chakri Sirindhorn. Clinical practice guideline for diabetes. Bangkok: The Endocrine Society of Thailand; 2017.
4. Anderson RJ FK, Clouse RE, Lustman PJ. The prevalence of comorbid depression in adults with diabetes: a meta-analysis. *Diabetes Care* 2001;24: 1069-78.
5. Maher D, Ford N, Unwin N. Priorities for developing countries in the global response to non-communicable diseases. *Global Health* 2012;8:14.
6. Wang J, He M, Zhao X. Depressive Symptoms, Family Functioning and Quality of Life in Chinese Patients with Type 2 Diabetes. *Can J Diabetes* 2015;39:507-12.
7. American Diabetes Association. Standards of medical care in diabetes-2016. *Diabetes Care* 2016;39 (Suppl 1): S13-22.
8. Reutrakul S, Deerochanawong C. Diabetes in Thailand: Status and Policy. *Curr Diab Rep* 2016;16:28.
9. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67: 361-70.
10. Nilchaikovit T, Lortrakul M, Phisansuthideth U. Hospital Anxiety and Depression Scale (Thai HADS). *J Psychiatr Assoc Thai* 1996;41:18-30.
11. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale. An updated literature review. *J Psychosom Res* 2002; 52:69-77.
12. Temrat C, Suwanmanee P. Health perception, and quality of service as perceived by persons receiving and annual health check up at Songklanagarind Hospital. *Songklanagarind J Nurs* 2016;36:148-58.

13. Wongwairoj A. Factors related to annual health check up of staff in life insurance company. Bangkok: Srinakharinwirot University; 2005.
14. Becker MH, Maiman LA. Sociobehavioral determinants of compliance with health and medical care recommendations. *Med Care* 1975;13:10-24.
15. Supphapiphon S, Buathong N, Suppapatiporn S. Reliability and validity of the family state and function assessment scale. *Chula Med J* 2019;63:[In press]
16. Epstein NB, Baldwin LM, Bishop DS. The McMaster family assessment device. *J Marital Fam Ther* 1983;9: 171-80.
17. Epstein NB, Bishop DS, Leven S. The McMaster model of family functioning. *J Marriage Fam Couns* 1978;4: 19-31.
18. Trangkasombat U. Family therapy and family counseling. Bangkok: Fuang Fa Printing; 1997.
19. Engum A, Mykletun A, Midthjell K, Holen A, Dahl AA. Depression and diabetes: a large population-based study of sociodemographic, lifestyle, and clinical factors associated with depression in type 1 and type 2 diabetes. *Diabetes care* 2005;28:1904-9.
20. Thaneerat T. Prevalence of depression, HbA [subscript 1]C level, and associated factors in outpatients with type 2 diabetes at endocrine clinic, King Chulalongkorn Memorial Hospital [thesis]. Bangkok: Chulalongkorn University; 2008.
21. Collins MM, Corcoran P, Perry IJ. Anxiety and depression symptoms in patients with diabetes. *Diabet Med* 2009;26:153-61.
22. Eashwar Anantha V, Gopalakrishnan S. Prevalence of depression in patients with type 2 diabetes mellitus and its association with inadequate glycemic. *Int J Community Med Public Health* 2017;4:3399-406.
23. AlBekairy A, AbuRuz S, Alsabani B, Alshehri A, Aldebasi T, Alkatheri A, et al. Exploring Factors Associated with Depression and Anxiety among Hospitalized Patients with Type 2 Diabetes Mellitus. *Med Princ Pract* 2017;26:547-53.
24. Kaur G, Tee GH, Ariaratnam S, Krishnapillai AS, China K. Depression, anxiety and stress symptoms among diabetics in Malaysia: a cross sectional study in an urban primary care setting. *BMC Fam Pract* 2013; 14:69.
25. Ali S, Stone MA, Peters JL, Davies MJ, Khunti K. The prevalence of co-morbid depression in adults with Type 2 diabetes: a systematic review and meta-analysis. *Diabet Med* 2006;23:1165-73.
26. Suppasso P. The prevalence of depression among type 2 diabetic patients in PhangKhon Hospital. *Srinagarind Med J* 2010;25:272-9.
27. Foran E, Hannigan A, Glynn L. Prevalence of depression in patients with type 2 diabetes mellitus in Irish primary care and the impact of depression on the control of diabetes. *Irish J Med Sci* 2015;184: 319-22.
28. Molee M. Self esteem, social support and self care behaviors of non insulin dependent diabetic patient in Chaophaya Abhai Bhubejhr Hospital, Prachinburi Province [thesis]. Bangkok: Chulalongkorn University; 2002.
29. Rungreangkulkij S. Prevalence and factors of depression among type 2 diabetic patients. *J Psychiatr Assoc Thai* 2014;59:287-98.
30. Azimi-Nezhad M, Ghayour-Mobarhan M, Parizadeh MR, Safarian M, Esmaceli H, Parizadeh SM, et al. Prevalence of type 2 diabetes mellitus in Iran and its relationship with gender, urbanisation, education, marital status and occupation. *Singapore Med J* 2008; 49:571-6.
31. Reiss S, Havercamp SM. Motivation in developmental context: A new method for studying self-actualization. *J Humanist Psychol* 2005;45:41-53.
32. Lakey B, Scoboria A. The relative contribution of trait and social influences to the links among perceived social support, affect, and self-esteem. *J Pers* 2005;73: 361-88.
33. Miller TA, Dimatteo MR. Importance of family/social support and impact on adherence to diabetic therapy. *Diabetes Metab Syndr Obes* 2013;6:421-6.
34. Azmoude E, Tafazoli M, Parnan A. Assessment of Family Functioning and Its Relationship to Quality of Life in Diabetic and Non-Diabetic Women. *J Caring Sci* 2016;5:231-9.
35. Takenaka H, Sato J, Suzuki T, Ban N. Family issues and family functioning of Japanese outpatients with type 2 diabetes: a cross-sectional study. *Biopsychosoc Med* 2013;7:13.
36. Sawatsri P. The use of herbs and indigenous food crop behaviors for blood glucose reduction among type 2 diabetic patients at subdistrict health promoting hospital, Ongkharak Hospital Network, Nakhonnayok Province. *J Phrapokkklao Nurs Coll* 2015;26:35-47.
37. Ghafarzagdegan R, Alizadeh Z, Abbasinya H, Malekhosseini A, Noruzi M, Matoury Poor P, et al. The effect of licorice on depression of nurses. *Complementary Med J of Faculty Nursing and Midwifery* 2016;6:1410-9.
38. Ofir R, Tamir S, Khatib S, Vaya J. Inhibition of serotonin re-uptake by licorice constituents. *J Mol Neurosci* 2003;20:135-40.