

## Original article

# Effects of general obesity on heart rate variability in Thai people with physical inactivity

Jantakan Saechee<sup>a</sup>, Kanita Wannasamai<sup>a</sup>, Pattharaporn Jaimanee<sup>a</sup>, Decha Pinkeaw<sup>b</sup>, Kroekkiat Chinda<sup>c</sup>, Tharnwimol Inthachai<sup>a,\*</sup>

<sup>a</sup>Department of Physical Therapy, Faculty of Allied Health Sciences, Naresuan University, Pitsanulok, Thailand

<sup>b</sup>Department of Physical Therapy, Faculty of Associated Medical Sciences, Chiang Mai University, Chiang Mai, Thailand

<sup>c</sup>Department of Physiology, Faculty of Medical Sciences, Naresuan University, Pitsanulok, Thailand

---

**Background:** Obesity is an important risk factor of cardiovascular diseases. Heart rate variability (HRV), and maximum oxygen consumption (VO<sub>2</sub>max) are parameters of cardiorespiratory function for determining aerobic capacity. However, the studies that investigated the effects of obesity on HRV, pulmonary function and VO<sub>2</sub> max in male participants aged 40 and 50 years, have been limited.

**Objectives:** This study aimed to compare HRV, pulmonary function and VO<sub>2</sub>max in male participants with normal body mass index (BMI) (18.5 – 22.9 kg/m<sup>2</sup>, n = 10), Class I obesity (25 - 29.9 kg/m<sup>2</sup>, n = 10) and also to use HRV to assess cardiopulmonary function in obese subjects.

**Methods:** In all, there were twenty healthy subjects, aged 40 - 50 years. Each subject was measured for their body composition, heart rate variability, pulmonary function and VO<sub>2</sub>max by cycle ergometer. The data were compared for the significant difference between groups by using unpaired *t*-test.

**Results:** The time domain parameters of HRV in the obese group were lower, low frequency and ratio of low and high frequency (LF/HF ratio) also were higher when compared with participants with normal BMI. Pulmonary function and maximum oxygen consumption were not significantly different between the groups (*P* > 0.05).

**Conclusion:** Autonomic balance was diminished in obese participants. Therefore, the assessment of HRV may be applied as a basic screening test before development of cardiovascular diseases.

**Keywords:** Obesity, heart rate variability, maximum oxygen consumption, pulmonary function, cardiovascular disease.

---

Currently, Thai society is changing from an agricultural culture to an industrial community. As such, people are living busier and more hurried lives, causing them to lose interest in taking care personal health causes many problems. <sup>(1)</sup> Obesity is a factor that causes people to be at higher risk of diseases and illness, all of which affect quality of life, society, and economy both within Thailand and around the world. <sup>(2)</sup>

Heart rate variability (HRV) is a parameter that evaluates the balance of autonomic nervous system (ANS) balance and predict the mortality and morbidity form cardiovascular diseases. <sup>(3)</sup> HRV comprises time domain parameters, the standard deviation of NN intervals (SDNN) and the root mean square of successive differences in NN intervals (RMSSD). Frequency domain of HRV consist of low frequency (LF), high frequency (HF), and LF/HF ratio indicating sympathetic and parasympathetic tone balance. <sup>(3)</sup> The research of Chintala and others which studied effects of being overweight had a depressed HRV value. <sup>(4)</sup> Additionally, visceral fat had a positive correlation with autonomic imbalance indicated by LF and LF/HF ratio. <sup>(5)</sup>

---

\*Correspondence to: Tharnwimol Inthachai, Department of Physical Therapy, Faculty of Allied Health Sciences, Naresuan University, Pitsanulok 65000, Thailand.

Email: tharnwimoli@nu.ac.th

Received: November 15, 2018

Revised: January 4, 2019

Accepted: February 4, 2019

VO<sub>2</sub> max is the ability of the cardiopulmonary system while doing daily physical activities.<sup>(6)</sup> Shanzia and others' findings about the effect of overweight and research done by Crump and others in obese participants found that VO<sub>2</sub> max was lower compared to normal BMI.<sup>(7,8)</sup> Obesity decreased the pulmonary function, leading to increase the work of breathing and need to exert more energy. As a result, obese people are likely to be more exhausted while performing daily physical activities.<sup>(9)</sup>

Even though there has been a number of studies about HRV, pulmonary function and VO<sub>2</sub> max in obese volunteers, the effects of overall obesity in middle-aged people in Thailand according to the Asian Pacific BMI criteria are still limited. In middle aged males, their metabolisms decrease due to chemicals and hormones changes, leading to raising the risk of being overweight or obese;<sup>(10)</sup> this largely results from improper health activity rather than from natural process of bodily change.<sup>(11)</sup> The incidence of overweight and obesity in Asia and Thailand, which found that the improper food consumption and sedentary lifestyle were important factors that cause people to be at risk of being overweight or obese.<sup>(12, 13)</sup>

The purpose of this study was to investigate the effects of overall sedentary obesity on HRV, pulmonary function, and VO<sub>2</sub> max and was also aimed to use ANS by measuring HRV in in male volunteers of age 40 - 50 years who are obese or having normal weight.

## Data collection procedures

### Participants

All participants were informed about the aims, consequences and they were free to make any decision to participate in the study according to the Declaration of Helsinki and was approved by the Naresuan University Institutional Review Board (IRB). Observational study in male subjects aged 40 - 50 years was done to the recruited 20 subjects by calculated form Chintala KK, *et al.*<sup>(4)</sup> Inclusion criteria were: nonsmoking male participants aged 40 - 50 years with normal BMI (18.5 - 22.9 kg/m<sup>2</sup>)<sup>(14)</sup>, Obese BMI (25 - 29.9 kg/m<sup>2</sup>)<sup>(14)</sup> and also had the level of healthy (20.5 - 27.4%, n = 10) and slightly high and higher (26.5 - 34.3%, , n = 10)<sup>(15)</sup> skinfold thickness. All subjects did a moderate or vigorous intensities of exercise less than 30 minutes for 5 days per week or 25 minutes 3 days per week.<sup>(16)</sup>

Informed consents were obtained from all subjects who could read and write Thai well before recruitment.

Exclusion criteria were as follows: people underlying cardiovascular, pulmonary and neurological diseases were excluded. People who had alcoholic beverages, caffeine or central nervous system (CNS) stimulation drugs within 48 hours and those had a musculoskeletal disease were also excluded.<sup>(17)</sup>

### Measurements of body composition

The volunteers were refrained from drinking alcoholic beverages, caffeine or CNS stimulation for at least 48 hours, and abstained exercise for at least 24 hours, had no food nor water for at least 4 hours, and abstained form diuretics for at least 7 days and also emptied their bladders for at least 30 minutes before the test.<sup>(18)</sup> After resting, all subjects were recorded their heart rate (HR) and blood pressure (BP) by cuff blood pressure (Omron HEM-7130, Omron Corporation, Japan) in the sitting position. The skinfold thicknesses were measured by skinfold caliper (Jama medical, model 5028, USA) with the intraclass correlation coefficient (ICC) value of 0.89 following as % BF formula = 0.39287 (sum of 3SKF) - 0.0105 (sum of 3SKF) 2 + 0.15772 (age) - 5.18845<sup>(15)</sup>

Body composition was measured by bioelectrical impedance analysis (BIA) (Omron HBF 375, Omron Corporation, Japan) with the intraclass correlation coefficient (ICC) 0.94 in the controlled the temperature room. All subjects were requested to clean their hands and feet before standing on the measuring unit with their arms raised and elbows extent straight until the device showed the results.<sup>(19)</sup>

### Pulmonary function test

All subjects followed the instructions and received the explanation and demonstration according to the standards of the American Thoracic Society (ATS) and the European Respiratory Society (ERS) criteria.<sup>(20)</sup> The normal values of lung function, forced vital capacity (FVC) and forced expiratory volume in one second (FEV1) and FEV1/ FVC ratio are more than 80% and >75% of the predictions, respectively. Pulmonary function test was performed through the mouthpiece and the device (Model ST90, Fukuda, Sangyo Co., Ltd. Japan) three times and brought the greatest value for interpretations.<sup>(20)</sup>

### Maximum oxygen consumption test

The researchers calculated VO<sub>2</sub> max by cycle

ergometer (*Monark*, Ergomedic 828 E, Sweden) following submaximal Astrand-Ryhming test protocol, with setting a rhythm at 50 repetitions per minutes (RPM). Sedentary male subjects were initially performed at 1 Kilo Pond intensity (300 - 450 kpm.min<sup>-1</sup>) to maintain HR between 120 - 170 beats/minutes.<sup>(21)</sup> HR for each stage was recorded throughout the test for six minutes. Constant HR at the fifth-sixth minutes was determined to correct VO<sub>2</sub> max. VO<sub>2</sub> max was calculated from their age factor and body weight to determined absolute (L/min) and relative VO<sub>2</sub> max (ml/kg/min), respectively.<sup>(21)</sup>

### Heart rate variability analysis

All subjects were instructed to breathe normally in supine position for at least 20 minutes and measured R-R interval from electrocardiogram using Power Lab and lab Chart V.8 (AD Instrument, Sydney, Australia) by recording electrode on the right arm (RA) and another on the left leg (LL) (lead II ECG).<sup>(22)</sup> The ECG activity was recorded continually with 200-Hz sampling frequency.<sup>(4)</sup> The R-R interval was measured in each average R-R interval was considered for the measurement after correction for artifacts and entropic beats.<sup>(23)</sup> Time domain analysis can be analyzed by calculation of indices based on R-R intervals.<sup>(23)</sup> Frequency domain calculations by spectral analysis of an array of R-R intervals was followed the Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology, LF and HF were ranged from 0.04 to 0.15 and 0.15 to 0.40 Hertz (Hz), respectively.<sup>(22)</sup>

### Statistical data analysis

Data were expressed as mean values ± standard deviation (SD). The normal distribution of data

was tested using Shapiro–Wilk tests. Unpaired *t*-tests were used to compare the differences between the two groups. The *P* < 0.05 was considered to be statistically significant difference.

## Results

### The general characteristic of the participants

The results of the study found that age and height of all subjects were no significant between the two groups (*P* > 0.05). However, the body weight, BMI and subcutaneous fat thickness were significantly different between the two groups. The mean BMI and the thickness of subcutaneous fat of the subjects with normal BMI were 21.83 ± 1.12 kg/m<sup>2</sup> and 23.85 ± 3.68 percentages. The body mass index group and the thickness of subcutaneous fat in the obesity group were 27.53 ± 1.61 kg/m<sup>2</sup> and 30.79 ± 2.11 percentages (Table 1).

There was significantly different in body fat mass between the groups. The body fat percentages of participants with normal and obese BMI were 17.68 ± 4.37 and 27.07 ± 3.26 percentages, respectively. There was no significant difference of muscle mass and visceral fat in both groups (*P* > 0.05). Heart rate, systolic and diastolic blood pressure showed no significant difference (*P* > 0.05) (Table 2).

### Comparison of pulmonary function and maximum oxygen consumption

There was no significant difference of pulmonary function and VO<sub>2</sub> max between the normal BMI and obesity groups (*P* > 0.05). The VO<sub>2</sub> max in normal BMI and obesity groups were 29.33 ± 3.29 and 26.41 ± 4.06 ml/kg/min, respectively, as shown in Table 3.

**Table 1.** General characteristics of the participants.

Parameters	Normal BMI	Obese BMI	<i>P</i> - value
Age (year)	44.70 ± 4.85	46.40 ± 5.10	0.286
Height (Centimeter)	167.30 ± 4.03	166.40 ± 4.79	0.511
Weight (Kilogram)	62.16 ± 4.76	76.18 ± 4.62	0.022
BMI (kg/m <sup>2</sup> ) (%Predict)	21.83 ± 1.12	27.53 ± 1.61	0.031
Skinfold thickness (%)	23.85 ± 3.68	30.79 ± 2.11	0.038
HR (BPM)	70.26 ± 8.06	68.16 ± 11.71	0.835
SBP (mmHg)	105.90 ± 7.77	110.80 ± 8.53	0.630
DBP (mmHg)	73.60 ± 6.31	75.30 ± 4.92	0.280

BMI = Body mass index, HR = Heart rate, BPM = Beat per minute, SBP = Systolic blood pressure, DBP = Diastolic blood pressure

**Table 2.** Bioelectrical body composition analysis.

Parameters	Normal BMI	Obese BMI	P - value
TBF (%)	17.68±4.37	27.07±3.26	0.038
VF (%)	4.25±1.75	6.71±2.81	0.897
FFM (%)	29.49±4.22	26.30±3.15	0.524

BMI = Body mass index, TBF = Total body fat, VF = Visceral fat body, FFM = Fat free mass

**Table 3.** Pulmonary function and maximum oxygen consumption.

Parameters	Normal BMI	Obese BMI	P - value
FEV1 (%Predict)	85.20±4.58	84.01±4.07	0.627
FVC (%Predict)	95.40±8.81	97.50±7.94	0.512
FEV1/FVC (%)	83.73±6.38	82.07±5.48	0.455
VO <sub>2</sub> max (ml/kg/min)	29.33±3.29	26.41±4.06	0.304

BMI = Body mass index, FEV1 = Forced expiratory volume in one second, FVC = Forced vital capacity, VO<sub>2</sub> max = Maximum oxygen consumption,

### Comparison of heart rate variability

Analysis of time domain parameters, SDNN and RMSSD of obese subjects at  $33.48 \pm 10.18$  and  $26.18 \pm 11.52$  milliseconds were significantly different, compared to volunteers with normal BMI at  $72.42 \pm 12.18$  and  $64.38 \pm 9.85$  milliseconds.

The frequency domain parameters, the LF and LF/HF ratio in both groups was different. The subjects with normal BMI had the LF and LF/HF values of  $49.69 \pm 13.38$  nu and  $0.95 \pm 0.17$ , respectively. Obese volunteers had LF and LF/HF values of  $76.02 \pm 8.56$  nu and  $2.05 \pm 0.75$ , respectively. There was no significant difference of HFnu between the two groups (Table 4)

### Discussion

In the present study, obese male subjects were found to have a decreased HRV, with both SDNN

and RMSSD parameters decreased as well as increased LF and LF/HF ratio, associated with increasing sympathetic activity, or autonomic imbalance.<sup>(24)</sup> There was no significant difference of pulmonary function and VO<sub>2</sub> max between groups, but VO<sub>2</sub> max was found to have a decreasing in obese participants.

Autonomic balance has an important role in regulating the cardiovascular function.<sup>(3)</sup> Obese subjects with higher fat mass in our study had a reduced SDNN and RMSSD as well as increased LF and LF/HF ratio, this suggested that body composition imbalance, increase of BMI and fat mass cause an autonomic dysfunction.<sup>(5)</sup> Obesity can lead to increased oxidative stress and inflammatory cytokine, associated with an increase in sympathetic activity as indicated by increased LF and LF/HF ratio which are factors of autonomic dysfunction.<sup>(25)</sup>

**Table 4.** Heart rate variability analysis.

Parameters	Normal BMI	Obese BMI	P - value
MeanRR (ms)	783.78±145.57	494.30±129.78	0.202
SDNN (ms)	72.42±12.18	33.48±10.18	0.027
RMSSD (ms)	64.38±9.85	26.18±11.52	0.031
LF (n.u.)	49.69±13.38	76.02±8.56	0.038
HF (n.u.)	60.38±10.41	45.56±11.85	0.187
LF/HF	0.95±0.17	2.05±0.75	0.024

BMI = Body mass index, Mean RR = Mean R to R interval, SDNN = Standard deviation of N to N interval, RMSSD = Root mean square of the successive differences, LF = Low frequency, HF = High frequency, ms = Millisecond

The cardiopulmonary system is influenced by the ANS and helps to accomplish daily routines efficiently, air that enters the lungs is transferred to disperse energy by the circulatory system for performing daily physical activity. <sup>(26)</sup> Previous studies showed that morbid obesity (indicated by a BMI greater than 30 kilograms per square meter) caused an increase of body fat percentage, especially in visceral body fat, and can lead to diminished chest and diaphragmatic movement and oxygen uptake <sup>(9, 27)</sup>, suggesting that muscle mass and visceral fat mass have a correlation with pulmonary function and  $VO_2$  max. <sup>(27, 28)</sup>

American College of Sports Medicine (ACSM) guidelines recommend a minimum of 30 minutes of moderately intense aerobic exercise for 5 days per week (150 minutes per week) or 25 minutes of vigorous exercise for 3 days per week (75 minutes per week) to improve cardiovascular endurance. <sup>(29)</sup> Overall obese middle aged males in our study had a higher body fat percentage all over their body, rather than extra fat limited to a specific region. <sup>(9)</sup> This results in an increase of total body fat, but visceral fat and muscle mass were not significantly different, compared to the subjects with normal BMI. This may have been a factor that caused a statistical difference of pulmonary and  $VO_2$  max. Furthermore, people with poor physical performance had a reduced HRV, and increased risk of cardiovascular diseases. <sup>(30)</sup> All subjects in our research do less exercise than the ACSM guidelines which suggested that increased BMI and fat mass might affect HRV and cause the decline of aerobic capacity in the future. <sup>(5)</sup>

Currently, HRV measurement has been a simple and reasonably-priced equipment, such as technological smart watches and phones, HRV can be applied to evaluate health promotion and sports competition planning. <sup>(24)</sup> The present study can serve as a path to health promotion for obese middle aged men, helping them to recognize their health through controlling BMI and body fat.

### Conclusion

In the present study, obese middle aged male had a depressed HRV. Therefore, control body composition and fat mass is not only essential to improve the ANS but also decrease the risk of cardiovascular diseases.

### Acknowledgements

The study is supported by grants from the Faculty

of Allied Health Sciences, Naresuan University, Thailand.

### Conflict of interest

The authors, hereby, declare no conflict of interest.

### References

1. Somrongthong R, Wongchalee S, Ramakrishnan C, Hongthong D, Yodmai K, Wongtongkam N. Influence of socioeconomic factors on daily life activities and quality of life of Thai elderly. *J Public Health Res* 2017; 6:862.
2. Kachur S, Lavie CJ, de Schutter A, Milani RV, Ventura HO. Obesity and cardiovascular diseases. *Minerva Med* 2017;108:212-28.
3. Verrier RL, Tan A. Heart rate, autonomic markers, and cardiac mortality. *Heart Rhythm* 2009;6(Suppl):S68-75.
4. Chintala KK, Krishna BH, NMR. Heart rate variability in overweight health care students: correlation with visceral fat. *J Clin Diagn Res* 2015;9:CC06-8.
5. Yadav RL, Yadav PK, Yadav LK, Agrawal K, Sah SK, Islam MN. Association between obesity and heart rate variability indices: an intuition toward cardiac autonomic alteration - a risk of CVD. *Diabetes Metab Syndr Obes* 2017;10:57-64.
6. Habibi E, Dehghan H, Moghiseh M, Hasanzadeh A. Study of the relationship between the aerobic capacity ( $VO_2$  max) and the rating of perceived exertion based on the measurement of heart beat in the metal industries Esfahan. *J Educ Health Promot* 2014;3:55.
7. Shazia SM, Badaam KM, Deore DN. Assessment of aerobic capacity in overweight young females: A cross-sectional study. *Int J Appl Basic Med Res* 2015; 5:18-20.
8. Crump C, Sundquist J, Winkleby MA, Sundquist K. Aerobic fitness, muscular strength and obesity in relation to risk of heart failure. *Heart* 2017;103:1780-7.
9. Mongkol S. The effect of obesity on respiratory and cardiovascular system. *J Assoc Med Sci* 2012;45:10-6.
10. Araujo AB, Wittert GA. Endocrinology of the aging male. *Best Pract Res Clin Endocrinol Metab* 2011;25: 303-19.
11. Leyk D, Ruther T, Wunderlich M, Sievert A, Essfeld D, Witzki A, et al. Physical performance in middle age and old age: good news for our sedentary and aging society. *Dtsch Arztebl Int* 2010;107:809-16.
12. Suppavitiporn S, Udompanyavit J, Suppavitiporn S. Overweight and obesity and the associated lifestyle factors in Thai adult individuals who came for service at King Chulalongkorn Memorial Hospital during

- August - September 2004. *Chula Med J* 2006;50:623-9.
13. Ramachandran A, Snehalatha C. Rising burden of obesity in Asia. *J Obes* 2010;2010.
  14. Stegenga H, Haines A, Jones K, Wilding J. Identification, assessment, and management of overweight and obesity: summary of updated NICE guidance. *BMJ* 2014;349:g6608.
  15. Samahito S. Assessment and the standard physical fitness for Thai people aged 19-59 years. Bangkok Sampachanya; 2013.
  16. Bennett JA, Winters-Stone K, Nail LM, Scherer J. Definitions of sedentary in physical-activity-intervention trials: a summary of the literature. *J Aging Phys Act* 2006;14:456-77.
  17. Myers J, Forman DE, Balady GJ, Franklin BA, Nelson-Worel J, Martin BJ, et al. Supervision of exercise testing by nonphysicians: a scientific statement from the American Heart Association. *Circulation* 2014;130:1014-27.
  18. Pereira C, Silva RAD, de Oliveira MR, Souza RDN, Borges RJ, Vieira ER. Effect of body mass index and fat mass on balance force platform measurements during a one-legged stance in older adults. *Aging Clin Exp Res* 2018;30:441-7.
  19. Oshima Y, Shiga T, Namba H, Kuno S. Estimation of whole-body skeletal muscle mass by bioelectrical impedance analysis in the standing position. *Obes Res Clin Pract* 2010;4:e1-e82.
  20. American Thoracic Society/European Respiratory Society. ATS/ERS Statement on respiratory muscle testing. *Am J Respir Crit Care Med* 2002;166:518-624.
  21. Cink RE, Thomas TR. Validity of the Astrand-Ryhming nomogram for predicting maximal oxygen intake. *Br J Sports Med* 1981;15:182-5.
  22. Triggiani AI, Valenzano A, Ciliberti MA, Moscatelli F, Villani S, Monda M, et al. Heart rate variability is reduced in underweight and overweight healthy adult women. *Clin Physiol Funct Imaging* 2017;37:162-7.
  23. Shaffer F, Ginsberg JP. An overview of heart rate variability metrics and norms. *Front Public Health* 2017;5:258.
  24. Manimmanakorn N, Manimmanakorn A, Vichiansiri R, Saengsuwan J, Naruemon L. Heart rate variability assessment and clinical uses. *J Thai Rehabil Med* 2018;28:32-6.
  25. Guarino D, Nannipieri M, Iervasi G, Taddei S, Bruno RM. The role of the autonomic nervous system in the pathophysiology of obesity. *Front Physiol* 2017;8:665.
  26. Hautala AJ, Kiviniemi AM, Tulppo MP. Individual responses to aerobic exercise: the role of the autonomic nervous system. *Neurosci Biobehav Rev* 2009;33:107-15.
  27. Littleton SW. Impact of obesity on respiratory function. *Respirology* 2012;17:43-9.
  28. Mondal H, Mishra SP. Effect of BMI, body fat percentage and fat free mass on maximal oxygen consumption in healthy young adults. *J Clin Diagn Res* 2017;11:CC17-CC20.
  29. Oja P, Titze S. Physical activity recommendations for public health: development and policy context. *EPMA J* 2011;2:253-9.
  30. Teisala T, Mutikainen S, Tolvanen A, Rottensteiner M, Leskinen T, Kaprio J, et al. Associations of physical activity, fitness, and body composition with heart rate variability-based indicators of stress and recovery on workdays: a cross-sectional study. *J Occup Med Toxicol* 2014;9:16.